

MEASUREMENT OF THE Q^2 -EVOLUTION OF THE BJORKEN INTEGRAL AT LOW Q^2

A. DEUR

*Thomas Jefferson National Accelerator Facility
12000 Jefferson Avenue
Newport News, VA 23606, USA
E-mail: deurpam@jlab.org*

We report on the extraction of the Q^2 -dependence of the Bjorken sum between $0.16 < Q^2 < 1.1 \text{ GeV}^2$. A twist analysis performed on these data shows that the higher twist corrections are small due to a cancellation between the twist-4 and 6 terms. The extraction of an effective strong coupling constant is discussed.

1. Bjorken Sum Rule

The Bjorken sum rule¹ played a central role in verifying in the spin sector that QCD is the correct gauge theory of strong interaction. It reads:

$$\int_0^1 (g_1^p - g_1^n) dx = \frac{g_a}{6} \left[1 - \frac{\alpha_s}{\pi} - 3.58 \left(\frac{\alpha_s}{\pi} \right)^2 - 20.21 \left(\frac{\alpha_s}{\pi} \right)^3 + \dots \right] + \sum_{i=1}^{\infty} \frac{\mu_{2i+2}^{p-n}}{Q^{2i}} \quad (1)$$

where the coefficients μ_{2i+2}^{p-n} are sums of elements of twist no higher than $2i$ and which Q^2 -dependence is given by DGLAP equations. The sum rule has been verified experimentally at $Q^2=5 \text{ GeV}^2$ to better than 10%.

Taking the $Q^2 \rightarrow 0$ limit relates the Bjorken sum rule to the Gerasimov-Drell-Hearn (GDH) sum rule² that stands at $Q^2 = 0$. This connection triggered the generalization of the GDH sum³ at finite Q^2 . Because the generalized GDH sum is in principle calculable at any Q^2 , it can help to study the transition from the hadronic to partonic degrees of freedom of strong interaction. However, the validity domains of the chiral perturbation theory (χ PT) at low Q^2 and pQCD calculations at higher Q^2 do not overlap. Lattice QCD should bridge the two domains but no calculation is available yet. The relation between the Bjorken and generalized GDH sum is:

$$\int_0^{1-} g_1^p - g_1^n dx = \frac{Q^2}{16\pi^2\alpha} (\text{GDH}^p(Q^2) - \text{GDH}^n(Q^2)) \quad (2)$$

Hence the Bjorken sum is essentially the $p - n$ flavor non-singlet part of the GDH sum, which yields simplifications: more reliable estimations of the unmeasured low- x part of the integral, simpler pQCD evolution equations and less complicated χ PT calculations. This might help in linking validity domains of pQCD and χ PT⁶. Hence the Bjorken sum appears as an important quantity to measure to understand the hadron-parton transition.

Precise data on the proton⁷, deuteron⁸, and ³He⁹ are available from the Thomas Jefferson National Accelerator Facility (JLab). We used them to extract the Bjorken sum from $Q^2 = 0.16$ to 1.1 GeV^2 . To combine proton and neutron data, the ³He data were reanalyzed at the same Q^2 as the proton data. For consistency, the unmeasured low- x part of the integral was re-evaluated for the three data sets using a consistent prescription. The results are shown on Fig. 1, left panel, together with SLAC E143 results in the resonance region¹⁰. The elastic contribution is not included. The systematic uncertainties are given by the horizontal bands. At low Q^2 χ PT calculations can be compared to the data. Calculation done in the heavy Baryon approximation¹² may be more reliable since they agree with data, up to about $Q^2 = 0.25 \text{ GeV}^2$ (to be compared to $Q^2 = 0.1 \text{ GeV}^2$ typically for singlet quantities). However, the results from Bernard *et al.*¹¹ which do not use this approximation, do not support this conclusion. The improved model of Soffer and Teryaev¹⁴ and the calculation from Burkert and Ioffe¹⁵ agree well with the data. The pQCD result at third order in α_s and leading twist is shown by the gray band. The width comes from the uncertainty on α_s . Somewhat surprisingly, the data agree with the leading twist calculation down to quite low Q^2 , which indicates that higher twist terms are small or cancel each others. We quantitatively addressed the question by performing a higher twist analysis¹³ that shows that the coefficient μ_4^{p-n} of the $1/Q^2$ correction has opposite sign and similar magnitude to the coefficient μ_6^{p-n} of the $1/Q^4$ correction at $Q^2 = 1. \text{ GeV}^2$: $\mu_6^{p-n}/Q^4 = 0.09 \pm 0.02$ and $\mu_4^{p-n}/Q^2 \simeq -0.06 \pm 0.02$. Higher Twist analyzes were also done separately on the proton¹⁶ and neutron¹⁷.

2. The Effective Strong Coupling Constant

The simple α_s -dependence of the Bjorken sum makes it an ideal tool to extract α_s . However, higher twists have to be known or negligible. This is not the case for us. This difficulty is avoided when using the concept of effective coupling constants¹⁸. Here, higher twists and QCD radiations at orders higher than α_s^2 are folded into α_s^{eff} . We obtain a quantity that is well defined at any Q^2 , well behaved when crossing Λ_{QCD} or a quark mass threshold

and that is renormalization scheme independent. However, α_s^{eff} becomes process dependent. This is not a problem since process-dependent coupling constants can be related using equations called “commensurate scale relations” that connect observables without scheme or scale ambiguity^{19,20}. The extracted α_s^{eff} is shown in Figure 1, right panel. Also plotted is the pQCD calculation of α_s^{eff} (light gray band), α_s calculated to order β_0 (pink band), α_s^{eff} calculated using the model of Burkert and Ioffe, and α_s^{eff} as extracted from SLAC E155 (open square). Satisfactorily, α_s^{eff} merges with α_s at large Q^2 as expected since their difference is due to pQCD radiative corrections and higher twists. At low Q^2 , the behavior of α_s^{eff} is constrained by the GDH sum rule. This, together with the steep rise α_s^{eff} at larger Q^2 strongly hint that α_s^{eff} has no significant scale dependence at low Q^2 . This possible “freezing” of α_s^{eff} at low Q^2 is a debated issue. Lower Q^2 data^{21,22} will have the definite word on this feature of α_s^{eff} .




Figure 1. Left: Q^2 -evolution of the Bjorken sum. The Bjorken sum formed using neutron data extracted from ^3He (D) data is shown by the triangles (squares). Right: Effective strong coupling constant.

3. Summary and Outlook

We have extracted the Bjorken sum in the Q^2 range of 0.16-1.1 GeV^2 . The parton to hadron gap, if smaller, is not bridged yet. The magnitudes of the higher twists were extracted. The higher twist effects appear to be small, due to a cancellation of the $1/Q^2$ and $1/Q^4$ terms. We extracted a physical coupling for the strong interaction. Data hint that α_s^{eff} loses its scale dependence at low Q^2 .

Acknowledgments

This work was supported by the U.S. Department of Energy (DOE) and the U.S. National Science Foundation. The Southeastern Universities Research Association operates the Thomas Jefferson National Accelerator Facility for the DOE under contract DE-AC05-84ER40150.

References

1. J. D. Bjorken, Phys. Rev. **148** 1467 (1966).
2. S. Gerasimov, Sov. J. Nucl. Phys. **2**, 1966. S. D. Drell and A. C. Hearn, Phys. Rev. Lett. **16**, 908 (1966).
3. M. Anselmino, B.L. Ioffe and E. Leader, Sov. J. Nucl. **49**, 136, (1989), X. Ji and J. Osborne, J.Phys. G27 127 (2001).
4. V. Bernard, T. R. Hemmert and Ulf-G. Meißner, Phys. Rev. D **67**, 076008 (2003), X. Ji, C. W. Kao and J. Osborne, Phys. Lett. **B472**, 1 (2000).
5. J. Ellis and R.L. Jaffe, Phys. Rev. **D9** 1444 (1974).
6. V. D. Burkert, Phys. Rev. D **63**, 097904 (2001).
7. CLAS collaboration: R. Fatemi *et al.*, Phys. Rev. Lett. **91**, 222002 (2003).
8. CLAS collaboration: J. Yun *et al.*, Phys. Rev. C **67**, 055204 (2003).
9. E94-010 collaboration: M. Amarian *et al.*, Phys. Rev. Lett. **89**, 242301 (2002); Phys. Rev. Lett. **92**, 022301 (2004).
10. K. Abe *et al.*, Phys. Rev. Lett. **78**, 815 (1997).
11. V. Bernard, T. R. Hemmert and Ulf-G. Meißner, Phys. Rev. D **67**, 076008 (2003).
12. X. Ji, C. W. Kao and J. Osborne, Phys. Lett. **B472**, 1 (2000).
13. A. Deur *et al.* hep-ex/0407007.
14. J. Soffer and O. V. Teryaev, hep-ph/041022
15. V. D. Burkert and B. L. Ioffe, Phys. Lett. **B296**, 223 (1992); J. Exp. Theor. Phys. **78**, 619 (1994).
16. M. Osipenko *et al.*, hep-ph/0404195.
17. E94-010 collaboration: Z. E. Meziani *et al.*, hep-ph/0404066.
18. G. Grunberg, Phys. Lett. **B95** 70 (1980); Phys. Rev. **D29** 2315 (1984)
19. S. J. Brodsky, H. J. Lu, Phys. Rev. **D51** 3652 (1995))
20. S.J. Brodsky, et al, Phys. Lett. **B372** 133 (1996)
21. J.P. Chen, A. Deur and F. Garibaldi, JLab experiment E97-110
22. M. Battaglieri, A. Deur, R. De Vita and M. Ripani, JLab experiment E03-006